# Raw Progesterone Female Health Supplements CAS 57-83-0 For Irregular Menstruation

# **Basic Information**

- Place of Origin:
- Minimum Order Quantity: 10Grams
- Price:
- Packaging Details:
- Delivery Time: 3-7days after received payment

China

USD

1kg/Foil Bag

5000KG Per Year

T/T, Western Union, PayPal

- Payment Terms:
- Supply Ability:



# **Product Specification**

- Product Name:
- Cas:
- Appearance:
- Purity:Usage:

• Highlight:

Irregular Menstruation Treatment

Progesterone

White Powder

57-83-0

99%

Progesterone Female Health Supplements, Female Health Supplements CAS 57-83-0



Our Product Introduction

## High Quality Raw Powder Progesterone for Irregular Menstruation Treatment CAS 57-83-0

### **Product introduction**

Product name	Progesterone
CAS No.	57-83-0
MF	C21H30O2
MW	314.46
EINECS No.	200-350-6
Density	d23 1.166; d20 1.171
Boiling Point	394.13°C (rough estimate)
Melting Point	128-132 °C (lit.)

### Description:

Progesterone (progesterone), also known as progesterone hormone, progesterone hormone, is the main biologically active progesterone secreted by the ovary, the molecular formula is C21H30O2. Before ovulation, the amount of progesterone hormone produced per day is 2 to 3 mg, mainly from the ovaries.

#### Uses:

- 1. Progesterone test.
- 2. Tire protection.
- 3. Dysfunctional uterine bleeding caused by anovulatory or luteal insufficiency.
- 4. Amenorrhea.
- 5. Dysmenorrhea.

 Menorrhagia.
For the treatment of menstrual disorders (such as amenorrhea and functional uterine bleeding), luteal insufficiency, threatened abortion and habitual abortion, and premenstrual syndrome.

8. As a sustained-release hormone drug in an intrauterine device.

#### Side Effects:

1. (1) Gastrointestinal reaction, poor appetite; (2) acne; (3) fluid retention and edema; (4) weight gain; (5) allergic dermatitis; (6) mental depression; (7) breast Pain; (8) Menstrual disorders, irregular bleeding or amenorrhea; (9) Long-term use can cause endometrial atrophy, decreased menstrual flow, and prone to vaginal fungal infection.

2. Rare adverse reactions include: headache; pain in the chest, buttocks, legs, especially the gastrocnemius muscle; weakness, numbness or pain in the arms and feet; sudden or unexplained shortness of breath, sudden slurred speech, sudden vision changes, relapse vision, blindness of varying degrees, etc.

3. Long-term application can cause abnormal liver function, increased incidence of ischemic heart disease; endometrial atrophy, decreased menstrual flow, and prone to vaginal fungal infection.

4. The application of some progestogens with high androgenic activity in early pregnancy may cause masculinization of female offspring; reproductive tract malformations in offspring, most commonly hypospadias.



